

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1642BJF

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 DEC 05 CASREACT(R) - Over 10 million reactions available
NEWS 4 DEC 14 2006 MeSH terms loaded in MEDLINE/LMEDLINE
NEWS 5 DEC 14 2006 MeSH terms loaded for MEDLINE file segment of TOXCENTER
NEWS 6 DEC 14 CA/CAPLUS to be enhanced with updated IPC codes
NEWS 7 DEC 21 IPC search and display fields enhanced in CA/CAPLUS with the
IPC reform
NEWS 8 DEC 23 New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/
USPAT2
NEWS 9 JAN 13 IPC 8 searching in IFIPAT, IFIUDB, and IFICDB
NEWS 10 JAN 13 New IPC 8 SEARCH, DISPLAY, and SELECT enhancements added to
INPADOC
NEWS 11 JAN 17 Pre-1988 INPI data added to MARPAT
NEWS 12 JAN 17 IPC 8 in the WPI family of databases including WPIFV
NEWS 13 JAN 30 Saved answer limit increased
NEWS 14 JAN 31 Monthly current-awareness alert (SDI) frequency
added to TULSA

NEWS EXPRESS JANUARY 03 CURRENT VERSION FOR WINDOWS IS V8.01,
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.
V8.0 USERS CAN OBTAIN THE UPGRADE TO V8.01 AT
<http://download.cas.org/express/v8.0-Discover/>

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 11:58:12 ON 10 FEB 2006

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CAPLUS' ENTERED AT 11:58:33 ON 10 FEB 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 10 Feb 2006 VOL 144 ISS 8
FILE LAST UPDATED: 9 Feb 2006 (20060209/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

```
=> s mc 192 or (MC192)
      33493 MC
      2040 MCS
      34960 MC
          (MC OR MCS)
      30687 192
          3 MC 192
          (MC(W)192)
          12 MC192
L1      15 MC 192 OR (MC192)

=> s cancer? or neoplas? or tumor
=> s cancer? or neoplas? or tumor?
      283169 CANCER?
      438119 NEOPLAS?
      417183 TUMOR?
L2      690968 CANCER? OR NEOPLAS? OR TUMOR?

=> s l1 and l2
L3      0 L1 AND L2

=> file pctfull
```

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	12.62	12.83

FILE 'PCTFULL' ENTERED AT 11:59:34 ON 10 FEB 2006
COPYRIGHT (C) 2006 Univentio

FILE LAST UPDATED: 3 JAN 2006 <20060103/UP>
MOST RECENT UPDATE WEEK: 200552 <200552/EW>
FILE COVERS 1978 TO DATE

>>> IMAGES ARE AVAILABLE ONLINE AND FOR EMAIL-PRINTS <<<

>>> UPDATING DELAYED DUE TO DELIVERY FORMAT CHANGES. <<<

>>> NEW IPC8 DATA AND FUNCTIONALITY NOT YET AVAILABLE IN THIS FILE.
USE IPC7 FORMAT FOR SEARCHING THE IPC. WATCH THIS SPACE FOR FURTHER

DEVELOPMENTS AND SEE OUR NEWS SECTION FOR FURTHER INFORMATION
ABOUT THE IPC REFORM <<<

```
=> s mc 192 or (MC192)
      32880 MC
      2921 MCS
      35224 MC
          (MC OR MCS)
      65659 192
          4 MC 192
          (MC(W)192)
          11 MC192
L4      14 MC 192 OR (MC192)

=> s cancer? or neoplas? or tumor?
      74539 CANCER?
      21534 NEOPLAS?
      62442 TUMOR?
L5      93014 CANCER? OR NEOPLAS? OR TUMOR?

=> s 14 and 15
L6      11 L4 AND L5

=> s conjugat? or coupl? or link? or attach?
      71814 CONJUGAT?
      314207 COUPL?
      285938 LINK?
      353436 ATTACH?
L7      617080 CONJUGAT? OR COUPL? OR LINK? OR ATTACH?

=> s 17 and 16
L8      11 L7 AND L6

=> s 18 not py>2000
      550224 PY>2000
L9      5 L8 NOT PY>2000

=> s 19 not py>1999
      630082 PY>1999
L10     4 L9 NOT PY>1999

=> s anticancer or (anti () cancer) or chemotherap
=> s anticancer or (anti () cancer) or chemotherap?
      13132 ANTICANCER
          9 ANTICANCERS
      13135 ANTICANCER
          (ANTICANCER OR ANTICANCERS)
      167501 ANTI
          165 ANTIS
      167532 ANTI
          (ANTI OR ANTIS)
          70375 CANCER
          27076 CANCERS
          72542 CANCER
          (CANCER OR CANCERS)
          10909 ANTI (W) CANCER
          29221 CHEMOTHERAP?
L11     38810 ANTICANCER OR (ANTI (W) CANCER) OR CHEMOTHERAP?

=> s 110 and 111
L12     1 L10 AND L11

=> d ibib
```

L12 ANSWER 1 OF 1 PCTFULL COPYRIGHT 2006 Univentio on STN
 ACCESSION NUMBER: 1997021732 PCTFULL ED 20020514
 TITLE (ENGLISH): DESIGN OF HORMONE-LIKE ANTIBODIES WITH AGONISTIC AND
 ANTAGONISTIC FUNCTIONS
 TITLE (FRENCH): OBTENTION D'ANTICORPS SIMULANT DES HORMONES ET AYANT
 DES PROPRIETES D'AGONISTE ET D'ANTAGONISTE
 INVENTOR(S): SARAGОВI, H., Uri;
 LeSAUTER, Lynne
 PATENT ASSIGNEE(S): MCGILL UNIVERSITY;
 SARAGОВI, H., Uri;
 LeSAUTER, Lynne
 LANGUAGE OF PUBL.: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

NUMBER	KIND	DATE
WO 9721732	A1	19970619

DESIGNATED STATES

W:

AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE
 ES FI GB GE HU IL IS JP KE KG KP KR KZ LC LK LR LS LT
 LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI
 SK TJ TM TR TT UA UG US UZ VN KE LS MW SD SZ UG AM AZ
 BY KG KZ MD RU TJ TM AT BE CH DE DK ES FI FR GB GR IE
 IT LU MC NL PT SE BF BJ CF CG CI CM GA GN ML MR NE SN
 TD TG

APPLICATION INFO.: WO 1996-CA815 A 19961206
 PRIORITY INFO.: GB 1995-9525180.7 19951208

=> d kwic

L12 ANSWER 1 OF 1 PCTFULL COPYRIGHT 2006 Univentio on STN
 ABEN . . . the NGF
 docking site. Such antibodies may be used for the treatment, diagnosis
 or prevention of neurological
 diseases, neuromas and neoplastic tumors which
 express TrkA receptors. Also these antibodies may be
 used to develop and screen for pharmaceutical agents which are
 agonistic. . .
 ABFR . . . des nerfs (FCN). Cet anticorps peut
 servir a diagnostiquer ou a soigner des maladies neurologiques, des
 nevromes et des tumeurs
 neoplasiques qui expriment les recepteurs TrkA. Egalement, ces
 anticorps peuvent etre utilises pour
 developper et evaluer les agents pharmaceutiques qui sont. . .
 DETD . . . NGF docking site. Such antibod-
 ies may be used for the diagnosis, treatment or pre-
 vention of neurological diseases, neuromas and neo-
 plastic tumors which express TrkA receptors. Also
 these antibodies may be used to develop and screen for
 pharmaceutical agents which are agonistic or antago-
 nistic. . .
 . . .
 vivo inhibition of nerve
 growth factor binding to TrkA receptor or the inter-
 nalization or downmodulation of the receptor, such as
 for inhibiting tumor growth in situ, for the treatment
 or prevention of neurological diseases, neuromas and
 neoplastic tumors which express TrkA receptors, for
 mapping hormone-receptor interactive sites and recep-
 tor domain-function correlation such as mapping TrkA
 docking sites, for screening pharmacological. . .

In accordance with the present invention there is also provided a method for the treatment of neurological diseases, neuromas and neoplastic tumors which

express TrkA receptors in a patient, which comprises administering an effective amount of an antibody of the present invention or a functional. . .

In accordance with the present invention there is also provided a pharmaceutical composition for the treatment of neurological diseases, neuromas and neoplastic tumors which express TrkA receptors, which comprises an effective amount of an antibody of the present invention or a functional fragment thereof in association. . .

In accordance with the present invention there is also provided a method for the prognosis or diagnosis of human tumors which comprises.

a) biopsy and immunocytochemistry of tumors using the antibody of the present invention and fragments thereof;

or

b) radiolabeling of the antibody of the present invention and fragments thereof and nuclear imaging. . .

In accordance with the present invention there is also provided a method for the treatment of human tumor of a patient which comprises the steps of.

a) coupling cytotoxic agents to the antibody of the present invention and fragments thereof;

b) administering the coupled antibody of step a) to the patient.

of the

central and/or peripheral nervous system, which comprises an effective amount of an antibody of the present invention or a functional fragment thereof coupled to a pharmaceutical agent in association with a pharmaceutically acceptable carrier. The pharmaceutical agent may be selected from the group consisting of radioligands,. . .

and

Scatchard plot analysis;

Fig. 6 illustrates the protection from apoptotic death by 5C3 and 5C3 Fabs;

Fig. 7 illustrates nuclear imaging of tumors in vivo with 5C3;

Fig. 8 illustrates the survival of TrkA-expressing cells in serum-free media by 5C3 and derivatives;

Fig. 9 illustrates the differentiation/neuritegenesis of human TrkA-expressing cells in serum media;

Fig. 10 illustrates Mab5C3 prevents TrkA-expressing tumor growth in vivo; and

Fig. 11 illustrates the topography of the CDRs of the present invention.

(Sigma, Saint Louis,

MO), anti-phosphotyrosine mAb 4G10 (UBI, Lake Placid,

NY), and anti-PI-3 kinase polyclonal serum (UBI) were purchased, mouse anti-rat p75 mAb MC192 ascites were a gift from P. Barker, and anti-p65 mAb 87 6 was grown in the laboratory.

products were characterized by SDS-PAGE under non-reducing or reducing conditions (100 mM 2-mercaptoethanol) to >98% purity. Control Fabs from anti-rat p75 mAb MC192 were similarly prepared.

NaCl, 0.5%
Tween™-20, pH 7.6) containing 1% BSA (Sigma), and immunoblotted with the indicated primary mAbs- Secondary antibodies were either horseradish peroxidase (HRP) conjugated goat anti-rabbit IgG (HRP-GaR), or goat anti-mouse IgG (HRP-G(xM) (Sigma). For detection the enhanced chemiluminescence (ECL) reagents (Amersham, Oakville, Ont.) were used following.

mAbs, mAb
5C3 Fab fragments, control mAb 192 Fab fragments or serum (final 5% FBS, normal growth conditions). where indicated, Fabs were externally cross-linked with goat anti-mouse Fab (GamFab, Sigma). Wells containing all culture conditions but no cells were used as blanks.

Monomeric 5C3 Fab protection was dose-dependent. However, equivalent or better protective effects were achieved when Fabs were externally cross-linked with G(xmFab antibodies. Specificity controls included those described in the previous section for whole mAb 5C3, plus 192 Fabs which had.

Aberrant expression of trkA mRNA and NGF responsiveness have been correlated with neurodegenerative disorders and neoplastic malignancy. Hence, TrkA-binding agents will be useful clinical tools in diagnosis, prognosis and perhaps treatment of these diseases. Indeed, mAb 5C3 binding is a positive prognostic marker for certain human neoplasias.

in neuroblastoma
Neuroblastomas* Number Positive Mixed Negative
Group 1 60 38 17 5
Group 2 53 1 3 5 35
*15 samples repeated after chemotherapy, at the time of second surgery or recurrence: 5C3 staining patterns remained unchanged in 14 tumors; 1 negative tumor sub-sequently positive post chemotherapy in regions of maturing elements.

- 25 -
Table 7
TrkA expression in other malignant tumors
Malignant tumor N=42 TrkA-Pos
Central nervous system tumors 6 0
Rhabdomyosarcomas 5 0
Primitive neuroectodermal tumors 6 0
Ewing's sarcomas 2 0
Wilm's tumors 6 1
Osteosarcomas 4 0
Melanomas 5 0

Breast carcinomas 5 0

Lung carcinomas 3 0

Table 8

TrkA detection by immunocytochemistry, RT-PCR
and western blot

IMMUNOCYTO. N. . .

Thus, artificial ligands of TrkA can induce receptor internalization and could be useful in delivering toxic agents to the cytoplasm of TrkA-expressing tumors.

TrkA-expressing neuronal 4 6 cells or fibroblastoid E25 cells undergo apoptotic death in serum free media but can be rescued. . . combined at suboptimal doses, as would be expected if mAb 5C3 bound and activated unoccupied TrkA receptors. Furthermore, morphological changes and increased attachment to plastic were observed in both the NGF and 5C3 treated cells.

Monomeric 5C3 Fabs protected E25 and 4 6 cells from apoptotic death. When Fabs were externally cross-linked using anti-Fab antibodies, a heightened response occurred. Since growth factor receptor activation requires bivalent binding, the monomeric 5C3 Fabs must have retained. . .

EXAMPLE I

PRELIMINARY STUDY OF THE EFFECT OF mAb 5C3 IN TUMOR GROWTH

Nude mice were injected subcutaneously (right abdominal side) with 2X10⁶ human TrkA expressing tumor cells. Two days post-injections tumors in all mice had begun to form. Mice were randomized prior to treatment. A total of four intraperitoneal injections of 100 micrograms. . .

The mAb 5C3 dramatically reduced the primary tumor weight with no observable metastatic invasion. A small fibrotic mass was localized at the site of injection in mAb 5C3 treated mice. In contrast, IgG treated mice had large, vascularized tumor masses, which metastasized to the liver, peritoneum gut and spleen. All animals had similar body weights ('30 grams).

Table 9

TREATMENT PRIMARY TUMOR WEIGHT METASTASIS WEIGHT (mg) mg)

5C3 mAb 50 ± 20 (fibrotic) NONE

mouse IgG 800 ± 250 350 ± 20

EXAMPLE II

The use of Mab 5C3 and its derivatives for the diagnosis, prognosis and localization of tumors. The in vivo targeting efficacy of agents that bind the NGF receptor p140 TrkA was evaluated.

Nuclear imaging studies were done after the injection of ^{99m}Tc-labeled compounds in nude mice bearing tumors. Kinetics of tumor targeting, blood clearance, and bioavailability were studied. Tumors that do not express TrkA were not targeted, demonstrating the

specificity in vivo. This biodistribution study demonstrates that receptor-specific molecule analogs may be useful and may be effective agents for the detection, diagnosis, and possible treatment of neoplasias involving overexpressed oncogenic receptors such as TrkA (Fig. 7).

99mTc-[SC3]
Ligand % id/g T/nT
tumor 1.25 1
blood 0.1 13
muscle 0.06 20
heart 0.10 13
lung 0.17 7.30
liver 0.61 2.10
spleen 0.13 9.42
kidney 1.48 0.9
large bowel 2.2 0.7

EXAMPLE III

5C3. . . relevant for binding TrkA. These were named CDR1, CDR2, and CDR3. Region CDR1 is connected to CDR2 by a 15 amino acid linker; CDR2 is connected to CDR3 by a 30 amino acid linker. Their secondary structures have been analyzed.

Monovalent Fabs of 5C3 obtained after papain digestion are also agonistic, especially when externally cross-linked by anti-Fabs. A smaller fragment of mAb 5C3 called CDR(R) also protects cells from apoptosis (Fig. 8).

than the 25 kDa NGF molecule and is still agonistic. CDR(R) is composed of 3 selected CDRs (out of 6 possible ones) linked by long spacer regions. Preliminary studies have suggested that actually only 2 of the 3 CDRs are relevant for binding to TrkA. Further, it is expected that even smaller fragments can be designed, e.g. upon removal of the linker regions.

CLMEN 5 The use of claim 3 or 4, for inhibiting tumor growth in situ.
6- The use of claim 3 or 4, for the treatment or prevention of neurological diseases, neuromas and neoplastic tumors which express TrkA receptors.

14 A method for the treatment of neurological diseases, neuromas and neoplastic tumors which express TrkA receptors in a patient, which comprises administering an effective amount of an antibody of claim 1 or a functional. . .

15 A pharmaceutical Composition for the treatment of neurological diseases, neuromas and neoplastic tumors which express TrkA receptors, which comprises an effective amount of an antibody of claim 1 or a functional fragment thereof in. . .

17 A method for the prognosis or diagnosis of human tumor which comprises:
a) biopsy and immunocytochemistry of tumors using the antibody of claim 12 and fragments thereof;
or

b) radiolabeling of the antibody of claim 12 and fragments thereof and nuclear imaging. . .

18. A method for the treatment of human tumor of a patient which comprises the steps of:

- a) coupling cytotoxic agents to the antibody of claim 12 and fragments thereof;
- b) administering the coupled antibody of step a) to said patient.

. . .
the central

and/or peripheral nervous system, which comprises an effective amount of an antibody of claim 1 or a functional fragment thereof coupled to a pharmaceutical

- 43 -

agent in association with a pharmaceutically acceptable carrier-

=> f his

101650 HIS

22 HISES

L13 101666 HIS

(HIS OR HISES)

=> file his

'HIS' IS NOT A VALID FILE NAME

SESSION CONTINUES IN FILE 'PCTFULL'

Enter "HELP FILE NAMES" at an arrow prompt (=>) for a list of files that are available. If you have requested multiple files, you can specify a corrected file name or you can enter "IGNORE" to continue accessing the remaining file names entered.

=> d his

(FILE 'HOME' ENTERED AT 11:58:12 ON 10 FEB 2006)

FILE 'CAPLUS' ENTERED AT 11:58:33 ON 10 FEB 2006

L1 15 S MC 192 OR (MC192)

L2 690968 S CANCER? OR NEOPLAS? OR TUMOR?

L3 0 S L1 AND L2

FILE 'PCTFULL' ENTERED AT 11:59:34 ON 10 FEB 2006

L4 14 S MC 192 OR (MC192)

L5 93014 S CANCER? OR NEOPLAS? OR TUMOR?

L6 11 S L4 AND L5

L7 617080 S CONJUGAT? OR COUPL? OR LINK? OR ATTACH?

L8 11 S L7 AND L6

L9 5 S L8 NOT PY>2000

L10 4 S L9 NOT PY>1999

L11 38810 S ANTICANCER OR (ANTI () CANCER) OR CHEMOTHERAP?

L12 1 S L10 AND L11

L13 101666 F HIS

=> s l12

L14 1 L10 AND L11

=> d ibib abs

L14 ANSWER 1 OF 1

ACCESSION NUMBER:

TITLE (ENGLISH):

TITLE (FRENCH):

PCTFULL COPYRIGHT 2006 Univentio on STN

1997021732 PCTFULL ED 20020514

DESIGN OF HORMONE-LIKE ANTIBODIES WITH AGONISTIC AND ANTAGONISTIC FUNCTIONS

OBTENTION D'ANTICORPS SIMULANT DES HORMONES ET AYANT

INVENTOR(S): DES PROPRIETES D'AGONISTE ET D'ANTAGONISTE
SARAGОВI, H., Uri;
LeSAUTER, Lynne
PATENT ASSIGNEE(S): MCGILL UNIVERSITY;
SARAGОВI, H., Uri;
LeSAUTER, Lynne
LANGUAGE OF PUBL.: English
DOCUMENT TYPE: Patent
PATENT INFORMATION:

NUMBER	KIND	DATE
WO 9721732	A1	19970619

DESIGNATED STATES

W:

AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE
ES FI GB GE HU IL IS JP KE KG KP KR KZ LC LK LR LS LT
LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI
SK TJ TM TR TT UA UG US UZ VN KE LS MW SD SZ UG AM AZ
BY KG KZ MD RU TJ TM AT BE CH DE DK ES FI FR GB GR IE
IT LU MC NL PT SE BF BJ CF CG CI CM GA GN ML MR NE SN
TD TG

APPLICATION INFO.: WO 1996-CA815 A 19961206
PRIORITY INFO.: GB 1995-9525180.7 19951208

ABEN The present invention relates to an agonistic anti-human TrkA mAb 5C3 which recognizes the NGF docking site. Such antibodies may be used for the treatment, diagnosis or prevention of neurological diseases, neuromas and neoplastic tumors which express TrkA receptors. Also these antibodies may be used to develop and screen for pharmaceutical agents which are agonistic or antagonistic to NGF binding to the TrkA receptors.

ABFR L'invention concerne un anticorps monoclonal (ACm) 5C3 dirige contre le recepteur TrkA humain et reconnaissant le site d'ancrage du facteur de croissance des nerfs (FCN). Cet anticorps peut servir a diagnostiquer ou a soigner des maladies neurologiques, des nevromes et des tumeurs neoplasiques qui expriment les recepteurs TrkA. Egalement, ces anticorps peuvent etre utilises pour developper et evaluer les agents pharmaceutiques qui sont des agonistes ou des antagonistes de la fixation du FCN au recepteur TrkA.

=> d 110 ibib 1-4

L10 ANSWER 1 OF 4 PCTFULL COPYRIGHT 2006 Univentio on STN
ACCESSION NUMBER: 1997021732 PCTFULL ED 20020514
TITLE (ENGLISH): DESIGN OF HORMONE-LIKE ANTIBODIES WITH AGONISTIC AND ANTAGONISTIC FUNCTIONS
TITLE (FRENCH): OBTENTION D'ANTICORPS SIMULANT DES HORMONES ET AYANT DES PROPRIETES D'AGONISTE ET D'ANTAGONISTE
INVENTOR(S): SARAGОВI, H., Uri;
LeSAUTER, Lynne
PATENT ASSIGNEE(S): MCGILL UNIVERSITY;
SARAGОВI, H., Uri;
LeSAUTER, Lynne
LANGUAGE OF PUBL.: English
DOCUMENT TYPE: Patent
PATENT INFORMATION:

NUMBER	KIND	DATE
WO 9721732	A1	19970619

DESIGNATED STATES

W: AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE
ES FI GB GE HU IL IS JP KE KG KP KR KZ LC LK LR LS LT
LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI
SK TJ TM TR TT UA UG US UZ VN KE LS MW SD SZ UG AM AZ
BY KG KZ MD RU TJ TM AT BE CH DE DK ES FI FR GB GR IE
IT LU MC NL PT SE BF BJ CF CG CI CM GA GN ML MR NE SN
TD TG

APPLICATION INFO.: WO 1996-CA815 A 19961206
PRIORITY INFO.: GB 1995-9525180.7 19951208

L10 ANSWER 2 OF 4 PCTFULL COPYRIGHT 2006 Univentio on STN
ACCESSION NUMBER: 1994020125 PCTFULL ED 20020513
TITLE (ENGLISH): TREATMENT OF MOTOR NEURON DISEASES WITH FIBROBLAST
GROWTH FACTOR-5 (FGF-5)
TITLE (FRENCH): TRAITEMENT DES AFFECTIONS DES NEURONES MOTEURS A L'AIDE
DU FACTEUR DE CROISSANCE DES FIBROBLASTES 5 (FGF-5)
INVENTOR(S): HUGHES, Richard, A.;
SENDTNER, Michael;
LINDHOLM, Dan;
THOENEN, Hans, F., E.
PATENT ASSIGNEE(S): MAX-PLANCK-GESELLSCHAFT ZUR FoERDERUNG DER
WISSENSCHAFTEN E.V.
LANGUAGE OF PUBL.: English
DOCUMENT TYPE: Patent
PATENT INFORMATION:

NUMBER	KIND	DATE

WO 9420125	A1	19940915

DESIGNATED STATES

W: AU CA CN CZ FI HU JP KR NO NZ PL RU SI SK UA AT BE CH
DE DK ES FR GB GR IE IT LU MC NL PT SE

APPLICATION INFO.: WO 1994-EP764 A 19940311
PRIORITY INFO.: US 1993-8/030,611 19930312

L10 ANSWER 3 OF 4 PCTFULL COPYRIGHT 2006 Univentio on STN
ACCESSION NUMBER: 1992009305 PCTFULL ED 20020513
TITLE (ENGLISH): ANTI CD-4 ANTIBODIES BLOCKING HIV-INDUCED SYNCYTIA
TITLE (FRENCH): ANTICORPS ANTI-CD4 BLOQUANT LES SYNCYTIA PROVOQUES PAR
LE VIH
INVENTOR(S): BURKLY, Linda, C.;
CHISHOLM, Patricia, L.;
THOMAS, David, W.;
ROSA, Margaret, D.;
ROSA, Joseph, J.
PATENT ASSIGNEE(S): BIOGEN, INC.;
BURKLY, Linda, C.;
CHISHOLM, Patricia, L.;
THOMAS, David, W.;
ROSA, Margaret, D.;
ROSA, Joseph, J.

LANGUAGE OF PUBL.: English
DOCUMENT TYPE: Patent
PATENT INFORMATION:

NUMBER	KIND	DATE

WO 9209305	A1	19920611

DESIGNATED STATES

W: AT AU BE BF BJ CA CF CG CH CI CM DE DK ES FR GA GB GN
GR IT JP LU ML MR NL SE SN TD TG US

APPLICATION INFO.: WO 1991-US8843 A 19911127
PRIORITY INFO.: US 1990-618,542 19901127

L10 ANSWER 4 OF 4 PCTFULL COPYRIGHT 2006 Univentio on STN
ACCESSION NUMBER: 1989001975 PCTFULL ED 20020513

TITLE (ENGLISH): RECOMBINANT ANTIBODIES AND METHODS FOR THEIR PRODUCTION
 TITLE (FRENCH): ANTICORPS RECOMBINANTS ET LEURS PROCEDES DE PRODUCTION
 INVENTOR(S): CATTANEO, Antonino;
 NEUBERGER, Michael, Samuel
 PATENT ASSIGNEE(S): CELLTECH LIMITED;
 CATTANEO, Antonino;
 NEUBERGER, Michael, Samuel
 LANGUAGE OF PUBL.: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

NUMBER	KIND	DATE
WO 8901975	A1	19890309

DESIGNATED STATES
 W: AT BE CH DE FR GB IT JP LU NL SE US
 APPLICATION INFO.: WO 1988-GB695 A 19880824
 PRIORITY INFO.: GB 1987-8719963 19870824

=> d 110 ibib kwic 4

L10 ANSWER 4 OF 4 PCTFULL COPYRIGHT 2006 Univentio on STN
 ACCESSION NUMBER: 1989001975 PCTFULL ED 20020513
 TITLE (ENGLISH): RECOMBINANT ANTIBODIES AND METHODS FOR THEIR PRODUCTION
 TITLE (FRENCH): ANTICORPS RECOMBINANTS ET LEURS PROCEDES DE PRODUCTION
 INVENTOR(S): CATTANEO, Antonino;
 NEUBERGER, Michael, Samuel
 PATENT ASSIGNEE(S): CELLTECH LIMITED;
 CATTANEO, Antonino;
 NEUBERGER, Michael, Samuel
 LANGUAGE OF PUBL.: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

NUMBER	KIND	DATE
WO 8901975	A1	19890309

DESIGNATED STATES
 W: AT BE CH DE FR GB IT JP LU NL SE US
 APPLICATION INFO.: WO 1988-GB695 A 19880824
 PRIORITY INFO.: GB 1987-8719963 19870824

DETD . . . as single
 units comprising two heavy and two light chains bound
 together in conventional fashion by disulphide.bonds.--
 Since IgG is monomeric, it readily attaches to its
 antigen, However, it is generally unable to activate
 any of the body's effector functions until it is
 aggregated.
 . . .
 derived from one species and a
 constant region derived from another species, altered
 antibodies, i.e. antibodies having a variable region
 from an Ig attached by peptide linkage to a
 co-expressed
 effector protein, and humanised antibodies, i.e.
 . . .
 use a glioma cell line as a host cell for
 the production of recombinant polymeric IgM. The glioma
 cell line is a cancerous cell line derived from glial
 cells from within the brain. Glial cells are classified
 as actively secreting cells. It is also possible. . .
 . . .
 or
 pancreatic cells, and neural cells, such as pheochromal

cells and, in particular, glial cells. Such actively secreting cells have the advantage that non-malignant (non-cancerous) cells can readily be cultured in vitro.

thus provide a host system for the production of recombinant polymeric IgM which will avoid any problems inherent in the use of cancerous host cells.

also contains a single intron between the regions encoding most of the leader peptide and the VH DJH region. The transcription unit is linked to a gene encoding guanidine phosphoribosyl transferase (gpt) which can be used as a selective marker.

cloned in a derivative of pSV2GPT (1) in which the BamHI site had been converted to a SacI site by use of linkers. The HS-V p 2 transcription unit was assembled in three parts. The promoter/transcription start region was obtained from plasmid pF1 (2) as. . .

HOPC 2020 but with the promoter/leader exon replaced by the hs-P70 promoter/VH-leader portion of plasmid pSV-HSVg2. The HSVX1 light chain transcription unit is linked to a gene endowing resistance to neomycin (neo) which can be used as a selective marker.

derivative of pSV2neo (6) in which the HindIII site has been destroyed by filling in and a new HindIII site created by linker insertion in the BamHI site. The promoter/transcription start/leader region is the same as in pSV-HSV W 2 except in that (i) the region. . .

(i) measuring receptors for nerve growth factor (NGF) (13); fluorescent staining with antibody MC 192 (14); immunoprecipitation of S100 protein (15); and immunofluorescent staining for glial fibrillar acidic protein (16). The PC12 cells were shown to respond to NGF.

Purification on hapten sorbents was carried out using NIP-caproate linked to Sepharose as previously described (9).

and analysed either after reduction (left panel) on a 7.5% SDS/polyacrylamide gel or unreduced (right panel) on a 4%polyacrylamide gel made using N,N'-diallyltartardiamide to cross-link. The positions of the originr IgMf IgE, IgG, and W markers are indicated.

```
=> file his
'HIS' IS NOT A VALID FILE NAME
SESSION CONTINUES IN FILE 'PCTFULL'
Enter "HELP FILE NAMES" at an arrow prompt (=>) for a list of files
that are available.  If you have requested multiple files, you can
specify a corrected file name or you can enter "IGNORE" to continue
accessing the remaining file names entered.
```

```
=> d his
```

(FILE 'HOME' ENTERED AT 11:58:12 ON 10 FEB 2006)

FILE 'CAPLUS' ENTERED AT 11:58:33 ON 10 FEB 2006

L1 15 S MC 192 OR (MC192)
L2 690968 S CANCER? OR NEOPLAS? OR TUMOR?
L3 0 S L1 AND L2

FILE 'PCTFULL' ENTERED AT 11:59:34 ON 10 FEB 2006

L4 14 S MC 192 OR (MC192)
L5 93014 S CANCER? OR NEOPLAS? OR TUMOR?
L6 11 S L4 AND L5
L7 617080 S CONJUGAT? OR COUPL? OR LINK? OR ATTACH?
L8 11 S L7 AND L6
L9 5 S L8 NOT PY>2000
L10 4 S L9 NOT PY>1999
L11 38810 S ANTICANCER OR (ANTI () CANCER) OR CHEMOTHERAP?
L12 1 S L10 AND L11
L13 101666 F HIS
L14 1 S L12

=> s l9 not l10

L15 1 L9 NOT L10

=> d ibib

L15 ANSWER 1 OF 1 PCTFULL COPYRIGHT 2006 Univentio on STN
ACCESSION NUMBER: 2000037103 PCTFULL ED 20020515
TITLE (ENGLISH): COMPOUNDS FOR INTRACELLULAR DELIVERY OF THERAPEUTIC
MOIETIES TO NERVE CELLS
TITLE (FRENCH): COMPOSES D'APPORT INTRACELLULAIRE DE GROUPES
CARACTERISTIQUES THERAPEUTIQUES A DES CELLULES
NERVEUSES
INVENTOR(S): WEBB, Robert, R.;
MCKEE, Constance, A.
PATENT ASSIGNEE(S): XAVOS;
WEBB, Robert, R.;
MCKEE, Constance, A.
LANGUAGE OF PUBL.: English
DOCUMENT TYPE: Patent
PATENT INFORMATION:

NUMBER	KIND	DATE

WO 2000037103	A2	20000629

DESIGNATED STATES

W:

AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE
DK DM EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE
KG KP KR KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX
NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT TZ UA
UG US UZ VN YU ZA ZW GH GM KE LS MW SD SL SZ TZ UG ZW
AM AZ BY KG KZ MD RU TJ TM AT BE CH CY DE DK ES FI FR
GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN GW
ML MR NE SN TD TG

APPLICATION INFO.: WO 1999-US28211 A 19991129
PRIORITY INFO.: US 1998-09/217,037 19981221

=> s IR3

L16 1062 IR3

=> s antibod?

L17 84196 ANTIBOD?

=> s l17 and l16

L18 191 L17 AND L16

=> d his

(FILE 'HOME' ENTERED AT 11:58:12 ON 10 FEB 2006)

FILE 'CAPLUS' ENTERED AT 11:58:33 ON 10 FEB 2006

L1 15 S MC 192 OR (MC192)
L2 690968 S CANCER? OR NEOPLAS? OR TUMOR?
L3 0 S L1 AND L2

FILE 'PCTFULL' ENTERED AT 11:59:34 ON 10 FEB 2006

L4 14 S MC 192 OR (MC192)
L5 93014 S CANCER? OR NEOPLAS? OR TUMOR?
L6 11 S L4 AND L5
L7 617080 S CONJUGAT? OR COUPL? OR LINK? OR ATTACH?
L8 11 S L7 AND L6
L9 5 S L8 NOT PY>2000
L10 4 S L9 NOT PY>1999
L11 38810 S ANTICANCER OR (ANTI () CANCER) OR CHEMOTHERAP?
L12 1 S L10 AND L11
L13 101666 F HIS
L14 1 S L12
L15 1 S L9 NOT L10
L16 1062 S IR3
L17 84196 S ANTIBOD?
L18 191 S L17 AND L16

=> s l18 and l5

L19 154 L18 AND L5

=> s l19 and l7

L20 152 L19 AND L7

=> s l11 and l20

L21 87 L11 AND L20

=> s l21 not py>2000

550224 PY>2000

L22 28 L21 NOT PY>2000

=> s l22 not py>1999

630082 PY>1999

L23 20 L22 NOT PY>1999

=> d ibib 1

L23 ANSWER 1 OF 20 PCTFULL COPYRIGHT 2006 Univentio on STN
ACCESSION NUMBER: 1999030727 PCTFULL ED 20020515
TITLE (ENGLISH): POLYMERIC PRODRUGS OF AMINO- AND HYDROXYL-CONTAINING
BIOACTIVE AGENTS
TITLE (FRENCH): PRODROGUES POLYMERIQUES D'AGENTS BIOACTIFS CONTENANT
AMINE OU HYDROXY
INVENTOR(S): GREENWALD, Richard, B.;
PENDRI, Annapurna;
CHOE, Yun, H.
PATENT ASSIGNEE(S): ENZON, INC.
LANGUAGE OF PUBL.: English
DOCUMENT TYPE: Patent
PATENT INFORMATION:

NUMBER	KIND	DATE

WO 9930727	A1	19990624

DESIGNATED STATES

W: AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE
 ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR
 KZ LC LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT
 RO RU SD SE SG SI SK SL TJ TM TR TT UA UG UZ VN YU ZW
 GH GM KE LS MW SD SZ UG ZW AM AZ BY KG KZ MD RU TJ TM
 AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE
 BF BJ CF CG CI CM GA GN GW ML MR NE SN TD TG

APPLICATION INFO.: WO 1998-US26565 A 19981214
 PRIORITY INFO.: US 1997-08/992,435 19971217
 US 1998-09/183,557 19981030

=> d kwic

L23 ANSWER 1 OF 20 PCTFULL COPYRIGHT 2006 Univentio on STN
 ABEN . . . to double prodrugs containing polymeric-based transport
 forms. These polymeric prodrugs are preferably of formula (I) wherein:
 L1 is a bifunctional linking
 moiety; in formula (a) B is H, a leaving group, a residue of an
 amine-containing moiety, or a
 residue of. . .

DETD . . . AND
 HYDROXYL-CONTAINING BIOACTIVE AGENTS
 TECHNICAL FIELD
 The present invention relates to double prodrugs. In particular, the
 invention relates to polymefic-based double prodrugs having reversible
 linkages
 involving amino and hydroxyl moieties of chemical compounds and
 biologically
 active materials such as enzymes, proteins and the like.
 . . .
 as with alkaloids, it has been
 determined that when only one or two polymers of less than about 10, 000
 daltons
 are conjugated thereto, the resulting conjugates are
 rapidly eliminated in vivo
 especially if a somewhat hydrolysis-resistant linkage is used.
 In fact, such
 0 conjugates are so rapidly cleared from the body that even if
 a hydrolysis-prone ester
 linkage is used, not enough of the parent molecule is
 regenerated in vivo. This is
 often not a concern with moieties such as proteins, enzymes and the like
 even when
 hydrolysis-resistant linkages are used. In those cases
 multiple polymer strands, each
 having a molecular weight of about 2-5 kDa, are used to further. . .

R14
 L, is a bifunctional linking moiety such as Y5
 7 R,
 or la
 f M-C
 R8 CFb P
 L Jn q
 Y,
 ll
 G is H or -C where
 B is H, a leaving. . .

. . .
 of a biologically active compound which remains
 after it has undergone a substitution reaction in which the prodrug
 carrier portion

has been attached.

capable of solubilizing amine-containing or hydroxyl-containing compounds and extending their half-life as compared to the native or even second prodrug counterparts. The linkage between the polymer 3 0 and the second prodrug compound as described above, hydrolyzes at a rate which 5 allows the compound to. . .

Methods of making and using the compounds and conjugates described herein are also provided.

R21 r S
9 Y4
RI
RI, Li]]C Y3 L; -y2'-G
f - I
RIO K4
M Ar V
IR t R51 U
wherein: R14
Li is a bifunctional linking moiety such as Y5
R15
7 R7 L Ja
or I
f M-C
R8 4CF42 P
L -J n q
Y,
G is H or where
B is H, . . .

C. LENKER MOIETY LI

As shown above, the invention includes bifunctional linking moiety L, which Y4 11 when combined with C, forms an amino acid residue linker, or when (p) is greater than one, a peptide residue linker.

I I

D. THE DOUBLE PRODRUG LINKAGE PORTION

Y4

The first labile bond of the double prodrug system, which joins the L, to) is selected to hydrolyze, such as via. . .

methyl or ethyl or substituted C,-6 alkyl. It is preferred that X is either 0 or NR,2-

2. Q Portion of the Linker

Alternatively, when L, includes Q, which is a moiety containing a fi-ee Y4

electron pair positioned three to six atoms from the], moiety, the polymer, RI,, is

preferably attached to Q via a heteroatom such as oxygen. In a preferred

12

embodiment, the free electron pair is five atoms from this. . .

In these embodiments, R, is attached to Q via NR₂, O, or S.
Thus, Q
assists hydrolysis of the prodrug linkage by anchimeric
assistance because the free
electron pair moiety can generate a three- to six-membered, but
preferably five-
membered, ring by-product upon hydrolysis of the preferably ester
linkage.

The linkages included in the compounds have hydrolysis rates
in the plasma
of the mammal being treated which is short enough to allow. . .

. . .
the disclosure of each is incorporated herein by reference. It will be
understood that the water-soluble polymer will be functionalized for
attachment to
the linkage via M, X or Q herein. As an example, the PEG
portion of the prodrugs
can be the following non-limiting compounds:. . .

In order to provide the desired hydrolyzable linkage, mono- or
di-acid
activated polymers such as PEG acids or PEG diacids can be used as well
as mono-
or di-PEG amines. . .

. . .
in the double prodrug must be sufficient so as to provide
sufficient circulation of the double prodrug before hydrolysis of the
linker.

Within the ranges provided above, polymers having molecular weight
ranges of at
least 20,000 are preferred in some aspects for chemotherapeutic
and organic
moieties. In the case of some nucleophiles such as certain proteins,
enzymes and
the like, polymers having a molecular weight. . .

. . .
form
(IV) with an activating moiety donor such as p-nitrophenyl chloride
(PNP-Cl)
(forming, for example, compound (V) in Figure 1); and optionally
e. attaching an amine-containing or hydroxyl-containing
compound
residue, e.g. the drug to be transported, to compound (V) by displacing
the leaving
group in a. . .

. . .
the first
method above and reacting it with an activating moiety donor such as
p-nitrophenyl
chloride (PNP-Cl) forming (VI) in Figure 1;
b. attaching an amine-containing or hydroxyl-containing
compound,
e.g. the drug to be transported, to the activated intermediate compound
(VI);
1 8
C. removing the protecting. . .

The resulting conjugated prodrug composition is then recovered
or isolated using
techniques known to those of ordinary skill, i.e. filtered,
recrystallized.

Once in place, the activated form of the PEG prodrug (or blocked prodrug) is ready for conjugation with an amine- or hydroxyl-containing compound.

Ara-C (cytosine arabinoside) and related anti-metabolite compounds, e.g., gemcitabine, etc. Alternatively, B can be a residue of an amine-containing cardiovascular agent, anti-neoplastic, anti-infective, anti-fungal such as nystatin and amphotericin B, anti-anxiety agent, gastrointestinal agent, central nervous system-activating agent, analgesic, fertility agent, contraceptive agent, anti-inflammatory.

Suitable proteins, polypeptides, enzymes, peptides and the like having at least one available amino group for polymer attachment include materials which have physiological or pharmacological activities as well as those which are able to catalyze reactions in organic solvents. The

factors and phospholipase-activating protein (PLAP). Other proteins of general biological or therapeutic interest include insulin, plant proteins such as lectins and ficins, tumor necrosis factors and related proteins, growth factors such as transforming growth factors, such as TGFa's or TGFP's and epidermal growth factors, hormones, 23 sornatomedins,

of a polypeptide demonstrating in vivo bioactivity. This includes amino acid sequences, nucleic acids (DNA, RNA) peptide 20 nucleic acids (PNA), antibody fragments, single chain binding proteins, see, for example U.S. Patent No. 4,946,778, disclosure of which is incorporated herein by reference, binding molecules including fusions of antibodies or fragments, polyclonal antibodies, monoclonal antibodies and catalytic antibodies.

herein is that there is available at least one (primary or secondary) amine 20 containing position which can react and link with a carrier portion and that there is not substantial loss of bioactivity after the double prodrug system releases and regenerates the

incorporation into the double prodrug compositions of the invention, may themselves be substances/compounds which are not active after hydrolytic release from the linked composition, but which will become active after undergoing a further chemical process/reaction. For example, an anticancer drug that is delivered to the bloodstream by the double

prodrug transport system, may remain inactive until entering a cancer or tumor cell, whereupon it is activated by the cancer or tumor cell chemistry, e.g., by an enzymatic reaction unique to that cell.

After conjugation, the remaining amine-containing compound is referred to as the residue of the unconjugated compound.

trees indigenous to China and *Nothapodytes foetida* trees indigenous to India. Camptothecin and related compounds and analogs are also known to be potential anticancer or antitumor agents and have been shown to exhibit these activities in vitro and in vivo. Camptothecin and related compounds are also.

The A ring can also be substituted in the 9-position with a straight or branched C1-30 alkyl or C1-47 alkoxy, optionally linked to the ring by a heteroatom i.e.- O or S. The B ring can be substituted in the 7-position with a.

OH moiety which is capable of reacting directly with activated forms of the polymer transport systems described herein or to the linking moiety intermediates, e.g. iminodiacetic acid, etc., which are then attached to a polymer such as PEG.

O

2 O OH

NH

Ph

ACO

5BZ

Paclitaxel: R¹, C61-15; R² CH₃CO; Taxotere: R¹, (CH₃)₃CO; R² H
These derivatives have been found to be effective anti-cancer agents.

parent compounds include, for example, certain low molecular weight biologically active proteins, enzymes and peptides, including peptido glycans, as well as other anti-tumor agents, cardiovascular agents such as forskolin; anti-neoplastics such as combretastatin, vinblastine, doxorubicin, Ara-C, maytansine, etc.; anti-infectives such as vancomycin, erythromycin, etc.; anti-fungals such as nystatin, amphotericin B, triazoles,

the double

2 O prodrug compositions of the invention, may themselves be substances/compounds

which are not active after hydrolytic release from the linked composition, but which

will become active after undergoing a further chemical process/reaction. For

example, an anticancer drug that is delivered to the bloodstream by the double

prodrug transport system, may remain inactive until entering a

cancer or tumor cell,
whereupon it is activated by the cancer or tumor
cell chemistry, e.g., by an
enzymatic reaction unique to that cell.

29

After conjugation, the remaining amine-or hydroxyl-containing
compound is
referred to as the residue of the unconjugated compound.

not only the reversible double prodrug system described
above but also a second polymeric transport system based on more
permanent types
of linkages. The hybrids can be prepared by at least two
methods. For example,
the benzyl-elimination-based double prodrug can be synthesized first and
then
PEGylated using any art-recognized activated polymer such as
thiazolidinyl thione-
or succinimidyl carbonate-activated PEG. Alternatively, the more
permanent

conjugation reaction can be performed first and the resultant
conjugates can be used
to form the double prodrug conjugates described herein. It
will be understood that
the hybrid systems will be better suited for proteins, enzymes and the
like where
multiple amino groups are available for attachment of the
polymeric transport
forms. For purposes of the present invention, activated polymers will be
understood to include polymers containing one or. . .

The activating terminal moiety can be any group which facilitates
conjugation of the polymers with the biologically active
material, i.e. protein,
enzyme, etc. either before or after the double prodrug transport system.

for, among other things, treating diseases which
are similar to those which are treated with the parent compound, e.g.
enzyme
replacement therapy, neoplastic disease, reducing
tumor burden, preventing
metastasis of neoplasms and preventing recurrences of
tumor/neoplastic growths in
mammals.

Synthesis of (24b): Compound 24b was prepared in a similar manner to
compound 24a using a 40 kDa MW PEG linker 4b in place of MW 5
kDa linker

4a. UV assay for this compound indicated the amount of daunorubicin
present is
2.3 %. In vitro and in vivo results for. . .

Synthesis of compound (26b): Compound 26b was prepared in a similar
manner

to compound 26a using a 40 kDa PEG linker 8b in place of the 5
kDaPEG linker

8a. UV assay for this compound indicated the amount of daunorubicin
present is
2.1 %.

Synthesis of compound (27b): Compound 27b was prepared in a similar
manner

to compound 27a using a 40 kDa PEG linker 6b in place of 5 kDaPEG linker 6a.

Example 29*

Synthesis of compound (27c): Compound 27c was prepared in a similar manner to compound 27a using a 40 kDa PEG linker 6c in place of 5 kDaPEG linker 6a.

Synthesis of compound (29b): Compound 29b was prepared in a similar manner to compound 29a using a 40 kDa PEG linker 14b in place of the 5 kDaPEG linker 14a. UV assay for this compound indicated the amount of daunorubicin present is 2.1%.

Synthesis of compound (31b): Compound 31b was prepared in a similar manner to compound 31a using a 40 kDa PEG linker 21b in place of the 5 kDa linker 21a.

Con-jugation of compound 2a or 32a to (L)-asparaginase: synthesis of compound (33): PEG linker 2a or 32a (450 mg, 0.084 mmol, 317 eq) was added to native (L)-asparaginase (37.5 mg, 416 gL, 0.00027 mmol) in. . . gentle stirring. The solution was stiffed at 30 °C for 30 minutes. A GPC column (Zorbax GF-450) was used to monitor PEG

conjugation: The PEG-Asp conjugate had a retention time of 8.5 min. At the end of the reaction (as evidenced by the absence of native enzyme),. . .

freshly prepared 33 was found to be 137 IU/mg (native asparaginase = 217 IU/mg). Protein modification of asparaginase with SS-PEG (a permanent linker) using a procedure corresponding to that described in the aforementioned U.S. Patent No. 4,179,337 gave a sin-fflar activity of 120 IU/mg. A. . .

Kinetics of hydrolysis of PEG conjugate of (L)-asparaginase (33) in rat plasma and buffer: The rate of hydrolysis of compound 33 in rat plasma was measured using a. . .

Synthesis of (34). a protein hybrid: con-jugation of (33) with SS-PEG (a permanent linker): PEG linker 2a (393 mg, 0.073 mmol, 70 eq) was reacted with native (L)-asparaginase (150 mg, 1.664 m.L, 0.00106 mmol) in 30 m.L. .

Demonstration of selective removal of reversible PEG linker (2a) from the hybrid (34): Generation of a permanently modified asparaffinasr,, compound (5, 100 mg of 34 is dissovied in 30. . . (Amicon) having a molecular weight cut off of 50,000 Daltons to remove free PEG which was formed by selective cleavage of the conjugates formed from the PEG-2a linker. The solution now contains only

SS-PEG conjugated asparaginase (35). Thus, the reversible linker is hydrolyzed, leaving only the relatively permanently bonded PEG attached to the asparaginase.

daunorubicin was given i.p. in balb/c mice bearing S.C. Madison 109 Lung Carcinoma at 1 & 4 days after inoculation. The median

tumor volume of treatment and control groups were measured and compared when the control group's median tumor volume reached approximately 2000 mm³.

was administered intravenously in nude mice bearing a human ovarian carcinoma xenografts at 1, 5 & 9 days after inoculation. The median tumor volume of treatment and control groups were measured and compared when the control group's median tumor volume reached approximately 1000 mm³.

200 William C. Rose. Evaluation of Madison 109 Lung Carcinoma as a Model for Screening Antitumor Drugs. Cancer Treatment Reports, 1981, 65, 299.

CLMEN. . . the formula:

I R₂₁ R₃₁ S

Y₄

R_i

R₁₁-L₁] C Y₃ C-Y₂'-G

R₁₀ K₄

L J_m Ar V

wherein: R t R₅₁ U

LI is a bifunctional linking moiety;

Y_J

G is H or -C-rs- where

B is H, a leaving group, a residue of an amine-containing moiety, or a residue of. . .

claim I 1, wherein A is selected from the group consisting of hydrogen, C₀₂₁₁, C₁₋₆ alkyl moieties, dialkyl acyl urea alkyls and IR₃]S I R₂]r

R_I Y₄ R_s

G'--Y₂-C Y₃ C

R₄ Ar R₁₀

M

61t

wherein G' is the same as G or another member of the group. . .

C Y₃ C N₂ B₂

I

Ar L M₄

IR R₅₁U

wherein M₂ is a cleavable or reversible protecting group;

LI is a bifunctional linking moiety; Y_J

B₂ is selected from the group consisting of H, UH] HU- and leaving groups;

Y, -4 are independently O, S, or NR₂

(r), . . .

step of

e. reacting the prodrug transport form of step d with an amine-containing or hydroxyl-containing compound residue to form a conjugate.

```

. . .
31S
Y
11 I1
M2 i]]C Y3 C_Y2 B2
I
Ar L K4
1R R51U
wherein M2 is a cleavable or reversible protecting group;
LI is a bifunctional linking moiety; YJ
11
B, is selected from the group consisting of H, OH, HC- and leaving
groups;
YI-4 are independently O or S or. . .

```

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	26.20	39.03

STN INTERNATIONAL LOGOFF AT 12:08:54 ON 10 FEB 2006

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1642BJF

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1		Web Page URLs for STN Seminar Schedule - N. America
NEWS 2		"Ask CAS" for self-help around the clock
NEWS 3	FEB 27	New STN AnaVist pricing effective March 1, 2006
NEWS 4	APR 04	STN AnaVist \$500 visualization usage credit offered
NEWS 5	MAY 10	CA/CAPLUS enhanced with 1900-1906 U.S. patent records
NEWS 6	MAY 11	KOREAPAT updates resume
NEWS 7	MAY 19	Derwent World Patents Index to be reloaded and enhanced
NEWS 8	MAY 30	IPC 8 Rolled-up Core codes added to CA/CAPLUS and USPATFULL/USPAT2
NEWS 9	MAY 30	The F-Term thesaurus is now available in CA/CAPLUS
NEWS 10	JUN 02	The first reclassification of IPC codes now complete in INPADOC
NEWS 11	JUN 26	TULSA/TULSA2 reloaded and enhanced with new search and